Colonoscopy Prep Instructions

Suprep

	Your procedure is scheduled on:
	Your arrival time is:
	(Your arrival time may change up to the day before the procedure)
	Location of procedure:
	☐ Tahoe Forest Hospital, Main Lobby Entrance 10121 Pine Avenue, Truckee, CA
	Incline Village Community Hospital880 Alder Avenue, Incline Village, NV
	An Endoscopy nurse will call you 3-7 days before your procedure to provide preparation information and to confirm your arrival time. Please be aware that arrival times may change up to a day before your procedure. If you miss our call, confirm your appointment with a voicemail at (530) 582-3455 or text (530) 562-7350.
	Post-Procedure Pick Up: Arrange for a responsible person to pick you up after your exam. Your driver must stay within 20 minutes of Tahoe Forest Hospital/Incline Village Community Hospital during the procedure. You may not drive yourself home or take TART buses for the remainder of the day. You may arrange for TART Connect if you live in Truckee or use UBER, if necessary.
1 \	WEEK BEFORE COLONOSCOPY
I	Purchase the following:
	Suprep: A prescription medicine available from your preferred pharmacy.
3 E	DAYS BEFORE COLONOSCOPY
ı	ow-Fiber Diet:
	eat a Low-Fiber Diet beginning 3 days prior to your exam, selecting low-fiber foods such as nilk, cheese, yogurt, chicken, fish, and eggs.
	Avoid high-fiber foods such as bread and cereal with seeds or nuts, raw vegetables, and ruits with peels and any berries.

one day before, and the 2nd part is taken the morning of your procedure.

1 DAY BEFORE COLONOSCOPY

When you wake up:

D

- Begin CLEAR LIQUID DIET only, including:
 - Water tap, sparkling, seltzer, coconut (no pulp)
 - **Juice** apple, white grape, white cranberry
 - Gatorade yellow, green, orange, clear
 - Jell-O yellow, green, orange

- Clear broth or bouillon chicken, beef, or veggie flavors
- Coffee, Tea without any creamer (dairy or non-dairy)
- Gummy Candies

At 5:00 pm the day before your colonoscopy:

— Start drinking your colonoscopy prep
Pour one (6 ounce) bottle of Suprep into the enclosed cup
Add cool drinking water to the 16 ounce line on cup and mix
Drink all liquid in cup
Drink two more 16 ounce containers of water over the next hour
— Please continue your clear liquid diet through the night to stay well hydrated
AY OF COLONOSCOPY
At (4 hours prior to arrival) drink the second part of your prep:
Pour one (6 ounce) bottle of Suprep into the enclosed cup
Add cool drinking water to the 16 ounce line on cup and mix
Drink all liquid in cup
Drink two more 16 ounce containers of water over the next hour
At (2 hrs prior to arrival) TAKE NOTHING BY MOUTH, INCLUDING WATER.
NO gum, candy, ice or chewing tobacco.
Take your prescription medications with a small sip of water.

Common Suprep Side Effects:

- Nausea, bloating, cramping, and vomiting may occur. If so, take a break from drinking the prep for 30 minutes. Resume drinking until it is all finished, taking breaks as needed.
- A bowel movement will usually occur within an hour after finishing prep and water, but can take 3-4 hours. Everyone is different.
- Bowel movements will become watery and frequent until colon fully cleanses. The result should be clear or pale yellow or pale green.